

## Nearest Points and Some Fixed Point Theorems for Weakly Compact Sets<sup>1</sup>

V. M. SEHGAL,\* S. P. SINGH,<sup>†</sup> AND R. E. SMITHSON\*

\* *Department of Mathematics, University of Wyoming,  
Laramie, Wyoming, 82071, and*

<sup>†</sup> *Department of Mathematics, Memorial University of Newfoundland,  
St. John's, Newfoundland, Canada, A1C 5S7*

*Submitted by V. Lakshmikantham*

*Received February 29, 1986*

We prove the following theorem and then derive several known results as corollaries. Let  $X$  be a convex subset of  $E$  and let  $f: (X, w) \rightarrow (E, T)$  be a continuous mapping. Let  $S$  be a nonempty, convex, and  $w$ -compact subset of  $X$  and  $K$  a  $w$ -compact subset of  $X$ . Let  $p \in P$  and let  $f$  satisfy the condition: for each  $y \in X \setminus K$ , there exists a  $x \in S$  such that  $p(x - fy) < p(y - fy)$ . Then there exists a  $u \in X$  satisfying  $p(u - fu) = \min\{p(x - fu): x \in X\}$ . © 1987 Academic Press, Inc.

Let  $(E, T)$  be a locally convex topological vector space with topology  $T$  and  $E^* = (E, T)^*$  be its topological dual. Let  $w = w(E, E^*)$  be the weak topology of  $E$  and let  $P$  and  $Q$  denote the family of continuous seminorms generating the topologies  $T$  and  $w$ , respectively. Clearly  $Q \subseteq P$  since  $w \subseteq T$ . We prove the following theorem and then derive several results as corollaries.

**THEOREM 1.** *Let  $X$  be a convex subset of  $E$  and let  $f: (X, w) \rightarrow (E, T)$  be a continuous mapping. Let  $S$  be a nonempty, convex, and  $w$ -compact subset of  $X$  and  $K$  a  $w$ -compact subset of  $X$ . Let  $p \in P$  and let  $f$  satisfy the condition: for each  $y \in X \setminus K$ , there exists a  $x \in S$  such that*

$$p(x - fy) < p(y - fy). \quad (1)$$

*Then there exists a  $u \in X$  satisfying*

$$p(u - fu) = \min\{p(x - fu): x \in X\}. \quad (2)$$

We first give some consequences of Theorem 1. Corollary 1 is a slight

<sup>1</sup> Research supported in part by NSERC Grant A5154.

extension of the result in [6] and it also improves a well-known result of Ky Fan [2] (see Corollary 2 below).

**COROLLARY 1.** *Let  $X$  be a nonempty convex and  $w$ -compact subset of  $E$  and let  $f: (X, w) \rightarrow (E, T)$  be a continuous mapping. Then for each  $p \in P$ , there exists a  $u \in X$  satisfying (2).*

*Proof.* Let  $X = K = S$  in Theorem 1. Since in this case  $X \setminus K = \emptyset$ , condition (1) is trivially satisfied. Consequently the result follows by Theorem 1.

*Remark.* It may be remarked that if  $E$  is additionally a Hausdorff space in Corollary 1, then it also follows that either  $f$  has a fixed point or there exist a  $p \in P$  and a  $u \in X$  with  $0 < p(u - fu) = \min\{p(x - fu): x \in X\}$ .

The proof is similar to Corollary 1 in [6].

**COROLLARY 2 (Ky Fan).** *Let  $X$  be a nonempty convex and compact subset of  $(E, T)$  and let  $f: (X, T) \rightarrow (E, T)$  be a continuous mapping. Then for each  $p \in P$ , there exists a  $u = u(p)$  satisfying (2).*

*Proof.* It suffices to show that  $f: (X, w) \rightarrow (E, T)$  is continuous. Let  $C$  be a  $T$ -closed subset of  $E$ . Then by hypothesis  $f^{-1}(C)$  is  $T$ -closed in  $X$  and hence is  $T$ -compact. This implies that  $f^{-1}(C)$  is  $w$ -compact and since  $w$ -topology is Hausdorff, it follows that  $f^{-1}(C)$  is  $w$ -closed.

Now we give the following lemma to be used in the proof of our theorem.

**LEMMA 1.** *Let  $X$  be a subset of  $E$  and let  $f: (X, w) \rightarrow (E, T)$  be a continuous mapping. Then for any  $p \in P$  and a fixed  $x \in E$ , the mapping  $g: X \rightarrow R$  (reals) defined by*

$$g(y) = p(y - fy) - p(x - fy) \quad (3)$$

*is  $w$ -lsc (lower semicontinuous).*

*Proof.* Note that the function  $g$  is lsc on  $(X, w)$  (see [4]) iff for each real  $r$ , the set  $A = \{y \in X: g(y) \leq r\}$  is  $w$ -closed. Let  $\{y_\alpha: \alpha \in A\}$  be a net in  $A$  and a  $y \in E$  such that  $y_\alpha \rightarrow y$  in  $w$ -topology. Then by hypothesis  $p(x - fy_\alpha) \rightarrow p(x - fy)$ . Choose a  $x^* \in E^*$  by the Hahn-Banach theorem (see [5]) such that  $x^*(y - fy) = p(y - fy)$  while  $x^*(z) \leq p(z)$  for each  $z \in E$ . Since  $y_\alpha - fy_\alpha \rightarrow y - fy$  in  $w$ -topology, it follows that

$$p(y - fy) - p(x - fy) = \lim_\alpha (x^*(y_\alpha - fy_\alpha) - p(x - fy_\alpha)). \quad (4)$$

However, for each  $\alpha \in A$ ,

$$x^*(y_\alpha - fy_\alpha) - p(x - fy_\alpha) \leq p(y_\alpha - fy_\alpha) - p(x - fy_\alpha) \leq r.$$

Consequently, it follows by (4) that

$$g(y) \leq r,$$

that is, the set  $A$  is  $w$ -closed.

The following result due to Allen [1] will be used in the proof of Theorem 1.

**THEOREM 2** (Allen [1]). *Let  $X$  be a nonempty convex set of a topological vector space  $E$  and let  $g$  be a real valued function defined on  $X \times X$  such that*

- (i) *for each fixed  $x \in X$ ,  $g(x, y)$  is lsc function of  $y$  on  $X$ ,*
- (ii) *for each fixed  $y \in X$ ,  $g(x, y)$  is quasi-concave function of  $x$  on  $X$ ,*
- (iii)  *$g(x, x) \leq 0$  for all  $x \in X$ .*
- (iv)  *$X$  has a nonempty compact, convex subset  $S$  such that  $\{y \in X: g(x, y) \leq 0 \text{ for all } x \in S\}$  is compact.*

Then there exists a point  $u \in X$  with  $g(x, u) \leq 0$  for all  $x \in X$ .

Recall that a real valued function  $f$  defined on a convex set  $X$  is quasi-concave iff for each real  $t$ , the set

$$\{x \in X: f(x) > t\} \quad \text{is convex.}$$

*Proof of Theorem 1.* Define a mapping  $G: X \times X \rightarrow R$  by

$$G(x, y) = p(y - fy) - p(x - fy).$$

We show that  $G$  satisfies the conditions of Theorem 2. Condition (1) follows by Lemma 1. Further, since for any  $y$ , and  $x_1, x_2 \in X$ ,  $a \geq 0$ ,  $b \geq 0$  with  $a + b = 1$ ,

$$p(ax_1 + bx_2 - fy) \leq ap(x_1 - fy) + bp(x_2 - fy),$$

it follows that for a fixed  $y \in X$ ,  $G(x, y)$  is a quasi-concave function of  $x$  on  $X$ . Condition (iii) holds trivially. To prove (iv) let

$$A = \{y \in X: G(x, y) \leq 0 \text{ for all } x \in S\}.$$

If  $A = \emptyset$ , then  $A$  is  $w$ -compact subset of  $S$ . If  $A \neq \emptyset$  and  $y \in A$ , then

$$p(y - fy) \leq p(x - fy) \quad \text{for all } x \in S.$$

In view of (1) this implies that  $y \in K$ . Thus  $A \subseteq K$ . Further,  $A$  is  $w$ -closed. In fact, for any fixed  $x \in S$ , by Lemma 1, we have

$$G(x, y) = g(y) = p(y - fy) - p(x - fy)$$

is  $w$ -lsc.

This implies that the set  $\{y: G(x, y) \leq 0\}$  is a  $w$ -closed subset of  $K$ . Consequently,

$$A = \bigcap_{x \in S} \{y: G(x, y) \leq 0\}$$

is  $w$ -closed subset of  $K$ . Since  $K$  is  $w$ -compact, it follows that  $A$  is  $w$ -compact. Thus, conditions of Theorem 2 are satisfied for  $(E, w)$ . Hence for some  $u \in X$ ,

$$p(u - fu) \leq p(x - fu)$$

for all  $x \in X$ .

In the setting of a normed linear space, we have the following.

**THEOREM 3.** *Let  $X$  be a convex subset of a normed linear space  $E$  and let  $f: (X, w) \rightarrow (E, \|\cdot\|)$  be a continuous mapping. Let  $S$  be a nonempty weakly compact, convex subset of  $X$ , and  $K$  a weakly compact subset of  $X$ . If  $f$  satisfies the condition: for each  $y \in X \setminus K$ , there exists a  $x \in S$  such that*

$$\|x - fy\| < \|y - fy\|$$

*then there exists a  $u \in X$  with  $\|u - fu\| = \min\{\|x - fu\|: x \in X\}$ .*

It may be pointed out that Theorem 3, and consequently Theorem 1, provides an extension of a recent result of Ky Fan (see [3, Theorem 7]).

## REFERENCES

1. G. ALLEN, Variational inequalities, complementarity problems, and duality theorems, *J. Math. Anal. Appl.* **58** (1977), 1-10.
2. KY FAN, Extension of two fixed point theorems of F. E. Browder, *Math. Z.* **112** (1969), 234-240.
3. KY FAN, Some properties of convex sets related to fixed point theorems, *Math. Ann.* **266** (1984), 519-537.
4. L. NACHBIN, "The Haar Integral," the University series, p. 24, Van Nostrand, New York, 1964.
5. A. P. ROBERTSON AND W. J. ROBERTSON, "Topological Vector Spaces," Cambridge Univ. Press, London, 1966.
6. V. M. SEHGAL AND S. P. SINGH, A variant of a fixed point theorem of Ky Fan, *Indian J. Math. (Silver Jubilee Year)* **25** (1983), 171-174.